

Comparison between Caribbean and eastern Pacific coral reefs

Jorge Cortés

CIMAR and Escuela de Biología, Universidad de Costa Rica, San Pedro, Costa Rica.

Resumen: El Caribe y el Pacífico Oriental están separados por una distancia corta en el tiempo y en el espacio, pero sus arrecifes coralinos son muy diferentes. Estas diferencias van más allá que sus características actuales, también difieren en su desarrollo Cuaternario. El crecimiento arrecifal en el Pacífico Oriental era escaso o limitado durante el Pleistoceno y Holoceno temprano debido a condiciones ambientales adversas. Además, eran más frágiles y expuestos a más bioerosión que los arrecifes del Caribe. La posibilidad de preservación de los arrecifes coralinos del Pacífico Oriental es comparativamente baja.

Key words: reef comparison, Caribbean, eastern Pacific, Quaternary.

The temporal and spatial distance separating the Caribbean and the eastern Pacific is short, as little as 50 km (Leschine 1981) and around 3.5 million years (Coates *et al.* 1992). This separation has resulted in very different oceanographic settings and coral reefs on both sides of the Central American isthmus (Glynn 1982). In this paper I extend Glynn's comparisons to include historical development of the reefs and more recent contrasting data from both sides.

Coral reef development in the Caribbean has been relatively constant during the whole of the Quaternary period, and fossil uplifted and submerged reefs are common in the region (Dodge *et al.* 1983, Macintyre 1988, Macintyre *et al.* 1991). In the eastern Pacific, uplifted reefs of Pleistocene or early Holocene age are not found (per. obs., based on field work and interviews with geologists that work in the area), there are no reports of submerged reefs, and there is only one example of an uplifted late Holocene reef (Urvina Bay, Galapagos: Colgan and Malmquist 1987), and of fossil corals (Sea of Cortez, Mexico: Squires 1959).

The absence of fossil reefs on the Pacific coast of the Americas may be due to several factors. It seems probable that no reef developed during the Pleistocene, because of extreme

changes in temperature, cooling due to intensified upwelling (Hays *et al.* 1989) and warming due to development of El Niño events (Colgan 1990). It was also a period of large shifts in sea level in a region with narrow continental shelves (Cortés 1986). During the late Holocene, reefs started to grow in the eastern Pacific in protected areas (Gulf of Chiriquí, Panamá: Glynn and Macintyre 1977; Golfo Dulce, Costa Rica: Cortés 1990a), but not in areas exposed to temperature extremes, where growth started less than 2,000 years ago (Caño and Cocos Islands, Costa Rica: Guzmán 1986, Macintyre *et al.* 1992; Galápagos Islands, Ecuador: Macintyre *et al.* 1992).

Additionally, recent research has shown that El Niño may kill whole reefs in the eastern Pacific (Glynn 1990) and that post-disturbance events may partially obliterate the reefs (Colgan 1990, Glynn and Colgan 1992, Guzmán and Cortés 1992). Furthermore, studies on siltation-stressed reefs in Golfo Dulce on the Pacific coast of Costa Rica, indicate that even after the corals have been killed, internal bioerosion by lithophagid bivalves continues (Cortés 1990b). Finally, monitoring of eastern Pacific reefs since the 1982-83 El Niño shows that recovery is slow (Glynn 1990, per. obs.).

The sluggish recovery of eastern Pacific reefs is exemplified by reefs killed over 350 years ago on the north Pacific coast of Costa Rica (Glynn *et al.* 1983) that have shown no signs of recovery.

Extensive death of coral communities due to some disturbance event, such as hurricanes (Woodley *et al.* 1981), post-event mortality due to predation (Knowlton *et al.* 1990) and slow recovery (Hughes 1992), have been observed in the Caribbean, but there are no reports of completely eroded reef frameworks. In addition, recovery of Caribbean reefs may be faster than eastern Pacific reefs because of more benign environmental conditions and/or more source populations supplying recruits in the Caribbean.

Several observations indicate that the possibility of preservation of eastern Pacific reefs is lower than for Caribbean reefs. Eastern Pacific reefs are smaller and more dispersed than their Caribbean counterparts (Glynn 1982). The inner structure of eastern Pacific reefs consists of loosely interlocking branching corals and/or unattached massive corals. Moreover, little or no submarine cementation is evident and crustose coralline algal pavements are weakly developed (Glynn and Macintyre 1977, Cortés 1990a). Also, bioerosion is probably higher in the eastern Pacific than in the Caribbean (Highsmith 1980). Caribbean reefs typically have a high degree of submarine cementation, abundant populations of crustose coralline algae, and massive reef structures (Goreau and Goreau 1973, Macintyre and Glynn 1976). Most eastern Pacific reef frameworks would probably break up and collapse during strong earthquakes. This does not seem to occur in the Caribbean. During a 7.5 earthquake on the Caribbean coast of Costa Rica, only a small section of a reef front collapsed, while the rest of the reef and all the other reef structures on the coast endured the movement (Cortés *et al.* 1992).

In conclusion, the differences between Caribbean and eastern Pacific coral reefs go beyond their present day characteristics (Glynn 1982). They also differ in their Quaternary development; most eastern Pacific coral reefs are late starters (growth began less than 2,000 years ago), and significant accretion is restricted to a few localities. In contrast, most shallow-water modern Caribbean reefs started to grow over 6,000 years ago and relict submerged reefs

document reef development throughout the Holocene transgression.

The absence of Quaternary fossil coral reefs in the eastern Pacific may be due to sparse or limited growth during the Pleistocene and early Holocene because of adverse environmental conditions. Also, because eastern Pacific coral reefs are more fragile than their Caribbean counterparts, the probability of them being preserved in the fossil record is relatively low.

ACKNOWLEDGEMENTS

Reviews of this paper by P.W. Glynn and I.G. Macintyre are greatly appreciated.

REFERENCES

- Coates, A.G., J.B.C. Jackson, L.S. Collins, T.M. Cronin, H.J. Dowsett, L.M. Bybell, P. Jung & J.A. Obando 1992. Closure of the Isthmus of Panama: the near-shore marine record in Costa Rica and western Panama. *Geol. Soc. Amer. Bull.* 104:814-828.
- Colgan, M.W. 1990. El Niño and the history of eastern Pacific reef building, p. 183-229. *In* P.W. Glynn (ed.) *Global ecological consequences of the 1982-83 El Niño-Southern Oscillation*. Elsevier, Amsterdam.
- Colgan, M.W. & D.L. Malmquist. 1987. The Urvin Bay uplift: a dry trek through a coral community. *Oceanogr. Mar. Biol.* 30:61-66.
- Cortés, J. 1986. Biogeografía de corales hermatípicos: el Istmo Centro Americano. *An. Inst. Cienc. Mar. Limnol. UNAM* 13:297-304.
- Cortés, J. 1990a. Coral reef decline in Golfo Dulce, Costa Rica, eastern Pacific: anthropogenic and natural disturbances. PhD dissertation, University of Miami, Coral Gables.
- Cortés, J. 1990b. The coral reefs of Golfo Dulce, Costa Rica: distribution and community structure. *Atoll Res. Bull.* 334:1-37.
- Cortés, J., R. Soto, C. Jiménez & A. Astorga. 1992. Earthquake associated mortality of intertidal and coral reef organisms (Caribbean of Costa Rica). *Proc. 7th Int. Coral Reef Symp.*, Guam: in press.
- Dodge, R.E., R.G. Fairbanks, L.K. Benninger & F. Murrase. 1983. Pleistocene sea levels from raised coral reefs of Haiti. *Science* 219:1423-1425.
- Glynn, P.W. 1982. Coral communities and their modifications relative to past and prospective Central American seaways. *Adv. Mar. Biol.* 19:91-132.

- Glynn, P.W. 1990. Coral mortality and disturbances to coral reefs in the tropical eastern Pacific, p. 55-126. *In* P.W. Glynn (ed.). Global ecological consequences of the 1982-83 El Niño-Southern Oscillation. Elsevier Oceanography Series, Amsterdam.
- Glynn, P.W. & M.W. Colgan. 1992. Sporadic disturbances in fluctuating coral reef environments: El Niño and coral reef development in the eastern Pacific. *Amer. Zool.* 32:707-718.
- Glynn, P.W. & I.G. Macintyre. 1977. Growth rate and age of coral reefs on the Pacific coast of Panamá. *Proc. 3rd Int. Coral Reef Symp.*, Miami 2:251-259.
- Glynn, P.W., E.M. Druffel & R.B. Dunbar. 1983. A dead Central American coral reef tract: possible link with the Little Ice Age. *J. Mar. Res.* 41:605-637.
- Goreau, T.F. & N.I. Goreau. 1973. The ecology of Jamaican coral reefs. II. Geomorphology, zonation, and sedimentary phases. *Bull. Mar. Sci.* 23:399-464.
- Guzmán, H.M. 1986. Estructura de la comunidad arrecifal de la Isla del Caño, Costa Rica, y el efecto de perturbaciones naturales severas. MSc. Thesis, Universidad de Costa Rica, San Pedro, Costa Rica.
- Guzmán, H.M. & J. Cortés. 1992. Cocos Island (Pacific of Costa Rica) coral reefs after the 1982-83 El Niño disturbance. *Rev. Biol. Trop.* 40: 309-324.
- Hays, P.E., N.G. Pisias & A.K. Roelofs. 1989. Paleooceanography of the eastern equatorial Pacific during the Pliocene: a high-resolution radiolarian study. *Paleoceanogr.* 4:57-73.
- Highsmith, R.C. 1980. Geographic patterns of coral bioerosion: a productivity hypothesis. *J. Exp. Mar. Biol. Ecol.* 46:177-196.
- Hughes, T. 1992. Long-term dynamics of Jamaican coral communities. *7th Int. Coral Reef Symp.*, Guam, 22-26 June 1992. Abstract p. 47.
- Knowlton, N., J.C. Lang & B.D. Keller. 1990. Case study of natural population collapse: post-hurricane predation on Jamaican staghorn corals. *Smiths. Contr. Mar. Sci.* 31:1-25.
- Leschine, T.M. 1981. The Panamanian sea-level canal. *Oceanus* 24:20-30.
- Macintyre, I. G. 1988. Modern coral reefs of western Atlantic: new geological perspective. *Amer. Ass. Petro. Geol. Bull.* 72:1360-1369.
- Macintyre, I.G. & P.W. Glynn. 1976. Evolution of modern Caribbean fringing reef, Galeta Point, Panama. *Amer. Ass. Petrol. Geol. Bull.* 60:1054-1072.
- Macintyre, I.G., P.W. Glynn & J. Cortés. 1992. Holocene reef history in the eastern Pacific: mainland Costa Rica, Caño Island, Cocos Island, and Galapagos Islands. *Proc. 7th Int. Coral Reef Symp.* Guam; in press.
- Macintyre, I.G., K. Rützler, J.N. Norris, K.P. Smith, S.D. Cairns, K.E. Bucker & R.S. Steneck. 1991. An early Holocene reef in the western Atlantic: submersible investigations of a dead relict reef off the west coast of Barbados, W.I. *Coral Reefs* 10:167-174.
- Squires, D.F. 1959. Results of the Puritan-American Museum of Natural History expedition to western Mexico. 7. Corals and coral reefs in the Gulf of California. *Bull. Amer. Mus. Nat. Hist.* 118:371-431.
- Woodley, J.D., E.A. Chornesky, P.A. Clifford, J.B.C. Jackson, L.S. Kaufman, N. Knowlton, J.C. Lang, M.P. Pearson, J.W. Porter, M.C. Rooney, K.W. Rylaarsdam, V.J. Tunnicliffe, C.M. Wahle, J.L. Wulff, A.S.G. Curtis, M.D. Dallmeyer, B.P. Jupp, M.A.R. Koehl, J. Neigel & E.M. Sides. 1981. Hurricane Allen's impact on Jamaican coral reefs. *Science* 214:749-755.